

Lista publikacji DNAPointer:

1) Juszcuk-Kubiak E, Flisikowski K, Wicińska K., Nucleotide sequence and variations of the bovine myocyte enhancer factor 2C (MEF2C) gene promoter in Bos Taurus cattle., Mol Biol Rep. 2010 Jun 29

<http://www.ncbi.nlm.nih.gov/pubmed/20585884>

2) Krawczyk A, Hasse-Lazar K, Pawlaczek A, Szpak-Ulczoek S, Krajewska J, Paliczka-Cieślak E, Jurecka-Lubieniecka B, Roskosz J, Chmielik E, Ziaja J, Cierpka L, Peczkowska M, Preibisz A, Januszewicz A, Otto M, Jarzab B., Germinal mutations of RET, SDHB, SDHD, and VHL genes in patients with apparently sporadic pheochromocytomas and paragangliomas, Endokrynol Pol. 2010 Jan-Feb;61(1):43-8.

<http://www.ncbi.nlm.nih.gov/pubmed/20205103>

3) Goluch D, Korwin-Kossakowska A, Prusak B, Pierzchala M, Urbanski P, Michaluk A, Sender G., The study of polymorphism within the promoter region of the osteopontin (OPN) gene in sows, Neuro Endocrinol Lett. 2009;30(4):525-9.

<http://www.ncbi.nlm.nih.gov/pubmed/20010486>

4) Szewczyk B, Barski P, Sihler W, Rabalski L, Skrzecz I, Hoyos-Carvajal L, de Souza ML., Detection and identification of baculovirus pesticides by multitemperature single-strand conformational polymorphism, J Environ Sci Health B. 2008 Sep;43(7):539-45.

<http://www.ncbi.nlm.nih.gov/pubmed/18803107>

5) Piekarska K, Jagielski M., Genetical conditioning of fluoroquinolone resistance mechanisms of clinical enterococcus faecalis strains. II. The mutations present in the qdrds of gyrA, gyrB, parC and parE genes, Med Dosw Mikrobiol. 2007;59(4):329-41.

<http://www.ncbi.nlm.nih.gov/pubmed/18416125>

6) Zasada AA, Jagielski M., Valuation for usefulness of selected chromosomal markers for Bacillus anthracis identification. II. Valuation for markers SSH and rpoB, Med Dosw Mikrobiol. 2006;58(4):355-62.

<http://www.ncbi.nlm.nih.gov/pubmed/17642313>

7) Borkowska E, Binka-Kowalska A, Constantinou M, Nawrocka A, Matych J, Kałuzewski B., P53 mutations in urinary bladder cancer patients from Central Poland., J Appl Genet. 2007;48(2):177-83.

<http://www.ncbi.nlm.nih.gov/pubmed/17495352>

8) Constantinou M, Binka-Kowalska A, Borkowska E, Zajac E, Jałmuzna P, Matych J, Nawrocka A, Kałuzewski B., Application of multiplex FISH, CGH and MSSCP techniques for cytogenetic and molecular analysis of transitional cell carcinoma (TCC) cells in voided urine specimens., J Appl Genet. 2006;47(3):273-5.

<http://www.ncbi.nlm.nih.gov/pubmed/16877809>

9) Korwin-Kossakowska A, Wszyńska-Koko J, Sender G., A novel polymorphism of the porcine prolactin gene (PRL)., Neuro Endocrinol Lett. 2006 Feb-Apr;27(1-2):241-6.

<http://www.ncbi.nlm.nih.gov/pubmed/16648817>

10) Bezak A, Kaczanowski R, Dossenbach-Glaninger A, Kucharczyk K, Lubitz W, Hopmeier P., Detection of single nucleotide polymorphisms in coagulation factor XI deficient patients by multitemperature single-strand conformation polymorphism analysis., J Clin Lab Anal. 2005;19(6):233-40.

<http://www.ncbi.nlm.nih.gov/pubmed/16302212>

11) Wszyńska-Koko J, Kurył J., A novel polymorphism in exon 1 of the porcine myogenin gene., J Appl Genet. 2005;46(4):399-402.

<http://www.ncbi.nlm.nih.gov/pubmed/16278514>

12) Ignacak M, Turek-Plewa J, Limon J, Trzeciak WH., A novel c.C2754 > T transition in the androgen receptor gene introduces the premature termination codon Q798X and results in a truncated form of the receptor., Gynecol Endocrinol. 2004 Oct;19(4):178-81.

<http://www.ncbi.nlm.nih.gov/pubmed/15724799>

13) Moric-Janiszewska E, Herbert E, Cholewa K, Filipecki A, Trusz-Gluza M, Wilczok T., Mutational screening of SCN5A linked disorders in Polish patients and their family members., J Appl Genet. 2004;45(3):383-90.

<http://www.ncbi.nlm.nih.gov/pubmed/15306732>

14) Gierczyński R, Jagielski M, Rastawicki W., Evaluation of the usefulness of multitemperature single-stranded conformational polymorphism (MSSCP) for differentiation of selected bacterial gene variants (alleles), Med Dosw Mikrobiol. 2003;55(2):147-55.

<http://www.ncbi.nlm.nih.gov/pubmed/14577194>

15) Rozycka A, Skorupska E, Kostyrko A, Trzeciak WH., Evidence for S284L mutation of the CHRNA4 in a white family with autosomal dominant nocturnal frontal lobe epilepsy., Epilepsia. 2003 Aug;44(8):1113-7.

<http://www.ncbi.nlm.nih.gov/pubmed/12887446>

16) Ignacak M, Niedziela M, Trzeciak WH., Transition C2718T in the AR gene, resulting in generation of a termination codon and truncated form of the androgen receptor, causes complete androgen insensitivity syndrome., J Appl Genet. 2002;43(1):109-14.

<http://www.ncbi.nlm.nih.gov/pubmed/12084976>

17) Moric E, Herbert E, Mazurek U, Samelska J, Cholewa K, Trusz-Gluza M, Wilczok T J., The KVLQT1 gene is not a common target for mutations in patients with various heart pathologies., Appl Genet. 2002;43(2):245-54.

<http://www.ncbi.nlm.nih.gov/pubmed/12080180>

18) Kaczanowski R, Trzeciak L, Kucharczyk K., Multitemperature single-strand conformation polymorphism., Electrophoresis. 2001 Oct;22(16):3539-45.

<http://www.ncbi.nlm.nih.gov/pubmed/11669539>